Spatial-Temporal Super-Resolution of SOHO and SDO Observations via Deep Learning

Chunhui Xu, Jason T. L. Wang, Haimin Wang, Yan Xu, Qin Li, Haodi Jiang, Jialiang Li, Vasyl Yurchyshyn Institute for Space Weather Sciences, New Jersey Institute of Technology, cx4@niit.edu

Overview: High-resolution magnetograms of solar active regions (ARs) are critical for studying fine-scale magnetic structures and their dynamic evolution, which are essential for understanding solar eruptions. While SOHO/MDI, a space-based instrument, provides continuous, full-disk magnetograms, its low spatial-temporal resolution limits its use. SDO/HMI, the successor of SOHO/MDI, provides higher spatial-temporal resolution observations. On the other hand, GST/NIRIS, a ground-based instrument with large apertures, provides much higher spatial resolutions, with limited FOVs and selected time windows. Here, we propose three deep learning models for super-resolution of solar ARs images: SolarCNN and GenMDI to enhance the spatial and temporal resolution, respectively, of line-of-sight (LOS) magnetograms of SOHO/MDI, and DeepHMI to improve LOS and transverse magnetograms of SDO/HMI guided by the observations

New Jersey Institute of Technology

1. SolarCNN: SolarCNN is an attention-aided convolutional neural network for spatially super-resolving solar active region (AR) magnetograms. It enhances SOHO/MDI data to match the quality of higher-resolution SDO/HMI observations.



2. Spatial Super-resolution of SOHO/MDI LOS Magnetogram Using SolarCNN: Case study on NOAA 11117 at 00:00:00 25 October 2010.



3. DeepHMI: DeepHMI is a conditional diffusion model for spatially super-resolving solar AR LOS magnetograms and transverse magnetograms. It enhances SDO/HMI data to match the quality of higher-resolution GST/NIRIS observations. The forward-diffusion process repeatedly adds Gaussian noise to a target image until pure noise is obtained. The model is trained with the backward-diffusion process. During testing, DeepHMI calculates a super-resolved result from a Gaussian noise.



4. Spatial Super-resolution of SDO/HMI LOS Magnetogram Using DeepHMI: Case study on NOAA 12371 at 16:48:00 UT on 22 June 2015.



5. GenMDI: GenMDI is a conditional diffusion model for temporally super-resolving solar AR LOS magnetograms. It enhances the temporal resolution of SOHO/MDI data from a 96-min cadence to a 12-min cadence.



6. Temporal Super-resolution of SOHO/MDI LOS Magnetogram Using GenMDI: Case study on NOAA 09802 from 17:36:00 UT to 19:12:00 UT on 04 February 2002.



Acknowledgement: This work was supported in part by U.S. NSF grants AGS-2149748, AGS-2228996, AGS-2229064, OAC-2320147, and NASA grants 80NSSC24K0548, 80NSSC24K0843, and 80NSSC24M0174.